

Temperature dependence of the magnetic hyperfine field at an s-p impurity diluted in $R\text{Ni}_2$

A. L. de Oliveira*

*Instituto Federal de Educação, Ciência e Tecnologia do Rio de Janeiro,
Campus Nilópolis, Rua Lúcio Tavares 1045, 26530-060, Nilópolis, RJ, Brazil*

C. M. Chaves† and A. Troper

Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud 150, 22290-180, Rio de Janeiro, RJ, Brazil

N. A. de Oliveira

*Instituto de Física Armando Dias Tavares, Universidade do Estado do Rio de Janeiro,
Rua São Francisco Xavier 524, 20550-013, Rio de Janeiro, Brazil*

We study the formation of local magnetic moments and magnetic hyperfine fields at an s - p impurity diluted in intermetallic Laves phases compounds $R\text{Ni}_2$ ($R = \text{Nd, Sm, Gd, Tb, Dy}$) at finite temperatures. We start with a clean host and later the impurity is introduced. The host has two-coupled (R and Ni) sublattice Hubbard Hamiltonians but the Ni sublattice can be disregarded because its d band, being full, is magnetically ineffective. Also, the effect of the $4f$ electrons of R is represented by a polarization of the d band that would be produced by the $4f$ magnetic field. This leaves us with a lattice of *effective* rare earth R -ions with only d electrons. For the dd electronic interaction we use the Hubbard-Stratonovich identity in a functional integral approach in the static saddle point approximation.

I. INTRODUCTION

The Laves phases intermetallic compounds $R\text{Ni}_2$ ($R =$ rare earth elements) crystallize in a cubic structure [1] and exhibit an interesting variety of behaviors related to the changes in their magnetic, electronic, and lattice structures. They exhibit magnetization associated both with the localized spins ($4f$) and with the itinerant electrons of the rare earth. Although the extensive studies made, the theoretical description of how some of their properties varies with temperature, due to the electronic correlations, still remains open. The Ni sublattice however can be disregarded because its d band, being full, is magnetically ineffective. Also the effect of the R $4f$ electrons is represented by a polarized d band produced by the $4f$ electrons magnetic field. This approximation generates an effective rare earth R lattice but whose interactions differ from the usual R pure metal because now the equivalent lattice constant is different (see the end of section III for a numerical indication of the difference between the two cases).

II. METHOD

II.1. The effective host

We start with a clean effective host described by the Hamiltonian of itinerant d -electrons:

$$\mathcal{H} = H_d^0 + H_d^1. \quad (1)$$

The first term of Eq.(1) is

$$H_d^0 = \sum_{l\sigma} \varepsilon_{0\sigma} d_{l\sigma}^\dagger d_{l\sigma} + \sum_{ll'\sigma} T_{ll'} d_{l\sigma}^\dagger d_{l'\sigma}, \quad (2)$$

where $\varepsilon_{0\sigma}$ is the energy of the center of the d band, now depending on the spin polarization; $d_{j\sigma}^\dagger$ ($d_{j\sigma}$) is creation (annihilation) operators, and $T_{ll'}$ is the hopping integrals between atoms from the R effective lattice.

H_d^1 represents the Coulomb interaction

$$H_d^1 = U \sum_l n_{l\uparrow} n_{l\downarrow}, \quad (3)$$

$n_{l\sigma}$ being the number operator.

The partition function \mathcal{Z} for the system described by Hamiltonian (1) can be written as

$$\mathcal{Z} = \int \prod_l d\nu_l d\xi_l \int \prod_j d\nu_j d\xi_j e^{-\beta \left[-\frac{U}{4} \sum_l (\nu_l^2 + \xi_l^2) \right]} \times \text{tr} e^{-\beta \left(H_d^0 - \frac{U}{2} \sum_{l\sigma} (i\nu_l + \sigma \xi_l) n_{l\sigma} \right)}. \quad (4)$$

where $\beta = 1/k_B T$, k_B is the Boltzman constant and T the temperature. The Hubbard-Stratonovich identity[2] was used in order to linearize the Coulomb interaction generating two floating fields, an electric, ν_l , and a magnetic, ξ_l . The static approximation has also been performed.

Now, because of the floating fields, the system, although pure, becomes disordered[3, 4]. In Eq.(4) we see that the floating fields create site dependent $\varepsilon_{l_0\sigma}$'s :

$$\varepsilon_{l_0\sigma} = \varepsilon_{0\sigma} - \frac{U}{2} (i\nu_l + \sigma \xi_l) \quad (5)$$

We then adopt the Coherent Potential Approximation[5, 6] (CPA) point of view in which

* alexandre.oliveira@ifrj.edu.br

† cmch@cbpf.br

the system is replaced by an ordered one with an uniform self energy Σ_σ in all sites $l \neq l_0$; in l_0 the energy $\varepsilon_{l_0\sigma}$ remains function of the fluctuation fields with respect to Σ_σ . The partition function in Eq (4) is also the partition function for the Hamiltonian below (which will be the one used to implement the CPA)

$$\tilde{H} = \tilde{H}^0 + \tilde{H}^1, \quad (6)$$

with

$$\tilde{H}^0 = \sum_\sigma [\varepsilon_{l_0\sigma} - \Sigma_\sigma] d_{l_0\sigma}^\dagger d_{l_0\sigma} = \sum_\sigma V_\sigma d_{l_0\sigma}^\dagger d_{l_0\sigma}, \quad (7)$$

and

$$\tilde{H}^1 = \sum_{l\sigma} \Sigma_\sigma d_{l\sigma}^\dagger d_{l\sigma} + \sum_{l'l'\sigma} T_{ll'} d_{l\sigma}^\dagger d_{l'\sigma} \quad (8)$$

In (8) both l and l' are $\neq l_0$. Summing over all possible l_0 we arrive at the self-consistency condition to determine Σ_σ :

$$\int d\xi_{l_0} d\nu_{l_0} \frac{V_\sigma(\xi_{l_0}\nu_{l_0})}{1 - V_\sigma(\xi_{l_0}\nu_{l_0}) g_{l_0l_0\sigma}(z)} P(\xi_{l_0}\nu_{l_0}) = 0. \quad (9)$$

where the probability distribution P , is

$$P(\xi_{l_0}\nu_{l_0}) = \frac{e^{-\beta\Psi(\xi_{l_0}\nu_{l_0})}}{\int d\xi_{l_0} d\nu_{l_0} e^{-\beta\Psi(\xi_{l_0}\nu_{l_0})}}, \quad (10)$$

and Ψ is the free energy associated to \tilde{H}^0 ,

$$\Psi(\xi_{l_0}\nu_{l_0}) = \frac{U}{4} (\xi_{l_0}^2 + \nu_{l_0}^2) + \frac{1}{\pi} \int d\varepsilon f(\varepsilon) \text{Im} \sum_\sigma \ln [1 - V_\sigma(\xi_{l_0}\nu_{l_0}) g_{l_0l_0\sigma}(z)], \quad (11)$$

$f(\varepsilon)$ is the Fermi function, $z = \varepsilon + i\delta$, $\delta \rightarrow 0^+$, and $g_{l_0l_0\sigma}(z)$ is the Green function for the hamiltonian \tilde{H}^0 .

Then, the partition function (4), in the CPA approach is reduced to

$$\mathcal{Z} = \int \prod_{l_0} d\nu_{l_0} d\xi_{l_0} e^{-\beta\Psi(\nu_{l_0}, \xi_{l_0})}. \quad (12)$$

II.2. The introduction of a s - p impurity

We now describe the effects caused by the introduction of a s - p impurity (say Cd). We add a potential $\mathcal{V}_{0\sigma}^{\text{imp}}$ to Equation 6,

$$\mathcal{V}_{0\sigma}^{\text{imp}} = [\varepsilon_\sigma^{\text{imp}} - \Sigma_\sigma(z)] d_{l_0\sigma}^\dagger d_{l_0\sigma}, \quad (13)$$

where the impurity energy $\varepsilon_\sigma^{\text{imp}}$ is self consistently determined using the Friedel [8] condition for the charges screening

$$\Delta Z = \Delta Z_\uparrow + \Delta Z_\downarrow, \quad (14)$$




FIG. 1. Local magnetic moment at Cd impurity diluted in $R\text{Ni}_2$ intermetallic host for light rare earths.

where ΔZ_σ is the total charge difference between the σ conduction electrons of the impurity and the host:

$$\Delta Z_\sigma = \ln \left\{ 1 - g_{l_0l_0\sigma}(\epsilon_F) [\varepsilon_\sigma^{\text{imp}} - \Sigma_\sigma(\epsilon_F)] \right\}. \quad (15)$$

Using the Dyson equation, the perturbed Green functions for this problem can be written as

$$G_{l_0l_0\sigma}(z) = \frac{g_{l_0l_0\sigma}(z)}{1 - g_{l_0l_0\sigma}(z) [\varepsilon_\sigma^{\text{imp}} - \Sigma_\sigma(z)]}. \quad (16)$$

The local density of states for the σ spin direction is

$$\rho_{0\sigma}(\varepsilon) = -\frac{1}{\pi} \text{Im} G_{l_0l_0\sigma}(z) \quad (17)$$

and the local occupation number is

$$n_{0\sigma} = \int_{-\infty}^{\epsilon_F} \rho_{0\sigma}(\varepsilon) f(\varepsilon) d\varepsilon. \quad (18)$$

So, the magnetic moment at the impurity site is

$$\tilde{m}(0) = \sum_\sigma \sigma n_{0\sigma} \quad (19)$$

and finally, we calculate the magnetic hyperfine field at the impurity site, assuming that it is proportional to $\tilde{m}(0)$, via the temperature independent $A(Z_{\text{imp}})$ Fermi-Segrè contact coupling parameter [7]:

$$B_{hf} = A(Z_{\text{imp}}) \tilde{m}(0), \quad (20)$$

III. RESULTS AND DISCUSSIONS

We have introduced an effective model to extend our previous zero temperature results [8] to investigate the




FIG. 2. Local magnetic moment at Cd impurity diluted in RNi_2 intermetallic host for heavy rare earths.

magnetic hyperfine fields at a s - p impurity such as Cd, in RNi_2 at finite temperatures. We adopt a standard

paramagnetic density of state extracted from first principle calculation [9]. For each RNi_2 compound our model has two adjustable parameters, namely $\varepsilon_{0\sigma}$ and U . These are determined by reproducing the zero temperature local magnetic moment and the critical temperature.

In Fig. 1, Fig. 2 and Fig. 3 we plot the calculated temperature dependence of the local magnetic moments for the light rare earth elements, for the heavy elements and the magnetic hyperfine fields at Cd as function of temperature. The results are in good agreement with the experimental results [10].

As stated before, in the effective R lattice the interactions are different from a R pure metal. From Fig. 2 we see that the local moment, in units of Bohr magneton μ_B , at $T = 0K$ is about 0.012 whereas in Ref. 11 it was found that in pure Gd metal it is about 0.05.

ACKNOWLEDGMENTS

We would like to acknowledge the support from the Brazilian agencies FAPERJ and CNPq.

-
- [1] K.H.J. Buschow. Rep. Prog. Phys., **40**, 1179, (1977).
 - [2] J. Hubbard, Phys. Rev. Lett. **3**, 77 (1959).
 - [3] H. Hasegawa, J. Phys. Soc. Japan **46**, 1504 (1979).
 - [4] Y. Takehashi, Phys. Rev B **41**, 9207 (1990).
 - [5] B. Velický, S. Kirkpatrick, and H. Ehrenreich, Phys. Rev. **175**, 747 (1968).
 - [6] H. Hasegawa, J. Phys. Soc. Jpn. **49**, 963 (1980).
 - [7] I.A. Campbell, J. Phys. C **12**, 1338 (1969).
 - [8] A.L. de Oliveira, N.A. de Oliveira and A. Troper, Phys. Rev. B **67**, 12411 (2003).
 - [9] H. Yamada, J. Inoue, K. Terao, S. Kanda and M. Shimizu, J. Phys. F: Met. Phys. **14**, 1943 (1984).
 - [10] S. Müller, P. de la Presa and M. Forker, Hyperfine Interac. **158**, 163 (2004).
 - [11] A. L. de Oliveira, M. V. Tovar Costa, N. A. de Oliveira and A. Troper, J. Appl. Phys. **81**, 4215 (1997).




FIG. 3. Magnetic hyperfine field at Cd impurity diluted in $R\text{Ni}_2$ intermetallic host. The experimental data were collected from Ref. 10.